

Plasma effects in neutrino-pair emission due to Cooper pairing of protons in superconducting neutron stars*

L. B. Leinson

Institute of Terrestrial Magnetism, Ionosphere and Radio Wave

Propagation RAS, 142190 Troitsk, Moscow Region, Russia

E-mail: leinson@izmiran.rssi.ru

Abstract

The neutrino emission due to formation and breaking of Cooper pairs of protons in superconducting cores of neutron stars is considered with taking into account the electromagnetic coupling of protons to ambient electrons. Our calculation shows that the contribution of the vector weak current to the $\nu\bar{\nu}$ emissivity of protons is much larger than that calculated by different authors without taking into account the plasma effects. Partial contribution of the pairing protons to the total neutrino radiation from the neutron star core is very sensitive to the critical temperatures for the proton and neutron pairing and can dominate in some domains of these parameters.

PACS number(s): 97.60.Jd, 95.30.Cq, 13.10.+q, 13.88.+e, 71.45.-d

Keywords: Neutron star, Neutrino radiation, Superconductivity, Plasma effects

Typeset using REVTeX

*Contribution to the International Workshop “HADRON PHYSICS 2002”, April 14 – 19, Bento Gonçalves, Rio Grande do Sul, Brazil.

When the temperature inside a neutron star core is lower than the critical temperature T_c for nucleon pairing, the nucleon matter consists of a condensate of Cooper pairs, which has thermal excitations in the form of not paired quasi-particles. Cooper-pair formation and pair-breaking coexist in statistical equilibrium and result in additional neutrino-pair emission from the neutron star. Under certain conditions, neutrino emission due to Cooper pairing of nucleons can dominate in neutrino energy losses from the neutron star. The mechanism of neutrino emission due to the singlet-state pairing of neutrons was proposed by Flowers et al. [1] many years ago.

Specifics of neutrino emission due to the proton pairing occurs because of a smallness of the vector weak coupling of a proton. Cooper pairing of protons takes place likely in 1S_0 -state [2]. When protons are treated non-relativistically, the total spin of the Cooper pair in the singlet-state is zero. By this reason, the axial-vector contribution of the proton weak current to the neutrino emissivity occurs only as a relativistic correction. Therefore, neutrino emission produced by the proton pairing is conventionally estimated negligible [3].

Such inference is made on the basis of calculations which ignored electromagnetic correlations among the charged particles in the QED plasma. Actually, protons are coupled to ambient electrons via the electromagnetic field. By undergoing a quantum transition to the paired state, protons polarize the medium, thus inducing the motion of electrons inside the Debye sphere around them. The electron weak current associated to this motion generates neutrinos coherently with the weak current of protons, because the wavelength λ of radiated neutrino pairs is much larger than the electron Debye screening distance D_e (typically, $D_e^2/\lambda^2 \sim 10^{-2}$). The induced radiation from ambient electrons many times exceeds the neutrino radiation from the initial proton pair [4], [5]. To demonstrate this we study the neutrino emissivity, caused by the proton pairing, by the use of the fluctuation-dissipation theorem. We consider the total energy which is emitted into neutrino pairs per unit volume and time. For one neutrino flavor, the emissivity is given by the following formula:

$$Q = \frac{G_F^2}{2} \int \frac{\omega}{\exp\left(\frac{\omega}{T}\right) - 1} 2 \operatorname{Im} \tilde{\Pi}_R^{\mu\nu}(K) \operatorname{Tr}(j_\mu j_\nu^*) \frac{d^3 k_1}{2\omega_1(2\pi)^3} \frac{d^3 k_2}{2\omega_2(2\pi)^3},$$

where the integration goes over the phase volume of neutrino and antineutrino of the total energy $\omega = \omega_1 + \omega_2$ and the total momentum $\mathbf{k} = \mathbf{k}_1 + \mathbf{k}_2$. In this formula G_F is the Fermi coupling constant, $\tilde{\Pi}^{\mu\nu}$ is the retarded polarization tensor of the medium which has ends at the weak vertex, and the neutrino weak current is of the standard form

$$j_\mu = \bar{\nu} \gamma_\mu (1 - \gamma_5) \nu. \quad (1)$$

To incorporate the collective plasma effects, one should take into account exchange of photons between charged particles in the plasma. This can be done in the Random phase approximation (RPA). In this case the weak polarization tensor of the medium is given by the diagrams shown in Fig. 1.

$$\Pi_V^{\text{RPA}} = C_V^2 \text{ (proton loop)} + c_V^2 \text{ (electron loop)} - \text{ (electron loop)} + C_V c_V \text{ (proton-electron loop)} - \text{ (electron loop)} + C_V c_V \text{ (electron loop)} - \text{ (proton loop)} + C_V^2 \text{ (proton loop)} - \text{ (proton loop)}$$

Fig. 1. RPA approximation to the weak polarization of the medium. Here $C_V = 0.04$ is the constant of the proton vector weak coupling with the neutrino field; c_V is the electron coupling constant, ($c_V = 0.96$ for electron neutrinos, and $c'_V = -0.04$ for muon and tau neutrinos).

Here the first term describes the contribution, which comes directly from the pairing protons, as shown in Fig. 2.

$$\text{ (proton loop)} = \text{ (proton loop with arrows)} + \text{ (proton loop with arrows)}$$

Fig. 2. Free-gas approximation to the weak polarization of protons.

The other terms are caused by the plasma polarization.

To demonstrate efficiency of the collective effects, the Q^{RPA} is plotted in Fig. 3 versus the dimensionless temperature $\tau = T/T_c$ together with that obtained in the free-gas approximation (FG) without collective effects. One can see that, the collective effects substantially enhance, the $\nu\bar{\nu}$ emissivity caused by the vector weak current of protons.




Fig. 3. Temperature dependence of the RPA vector weak current contribution to the neutrino emissivity in comparison with that obtained in the free-gas approximation (FG). The emissivities are shown versus the dimensionless temperature $\tau = T/T_c$ for beta-equilibrium

nuclear matter.

The effect of neutron superfluidity and/or proton superconductivity on different neutrino reactions in the neutron star core is very complicated. Different neutrino production mechanisms can dominate at different cooling stages depending on the temperature, and the matter density, and vary along with the chosen parameters T_{cp} , T_{cn} , which are the critical temperatures for the proton and neutron pairing. The Fig. 4 shows efficiency of different reactions at the baryon density $\rho = 2\rho_0$, where the direct Urca process is forbidden. Neutrino emissivities due to the singlet-state proton pairing, and the triplet-state neutron pairing are plotted in logarithmic scale against the temperature together with the total emissivity of two branches of the modified Urca processes, and the total bremsstrahlung emissivity caused by nn-, np-, and pp-scattering, which are suppressed due to the neutron superfluidity and/or proton superconductivity. Two panels of Fig. 4 differ only by different choice of parameters T_{cp} and T_{cn} . On the left panel we took $T_{cp} = 5.6 \times 10^8$ K, and $T_{cn} = 5.6 \times 10^9$ K. On the right panel we assume $T_{cp} = 3.5 \times 10^9$ K, and $T_{cn} = 8.5 \times 10^8$ K. Partial contributions of the proton and neutron pairing to the total energy losses are very sensitive to the corresponding critical temperatures.




Fig. 4. Temperature dependence of the neutrino emissivity in different reactions for β -equilibrium nuclear matter of the density $\rho = 2\rho_0$. The neutrino emissivity due to the triplet-state neutron pairing is shown by dashed line. The solid line is the neutrino emissivity due to the singlet-state proton pairing. Dot-and-dash line shows the total bremsstrahlung emissivity caused by nn-, np-, and pp-scattering; the dotted line exhibits the total emissivity of two branches of the modified Urca processes. On the left panel we took $T_{cp} = 5.6 \times 10^8$ K, and $T_{cn} = 5.6 \times 10^9$ K. On the right panel we assume $T_{cp} = 3.5 \times 10^9$ K, and $T_{cn} = 8.5 \times 10^8$ K.

REFERENCES

- [1] E. Flowers, M. Ruderman, P. Sutherland, ApJ, **205** (1976) 541.
- [2] T. Takatsuka, R. Tamagaki, Progr. Theor. Phys. Suppl. , **112** (1993) 27.
- [3] A. D. Kaminker, P. Haensel, and D. G. Yakovlev, A&A 345 (1999) L14.
- [4] L. B. Leinson, Phys. Lett., **B473** (2000) 318
- [5] L. B. Leinson, Nucl. Phys., **A687** (2001) 489.